

Potential Threat of Rabies Virus from Bat Bite in Nepal

Yogendra Shah^{1,2*}, Dhan Kumar Pant², Kishor Pandey^{1,3}, Ajay Poudel⁴, Bimal Dahal⁵, Krishna Prasad Pant^{1,6}, Basu Dev Pandey^{1,7}

¹Everest International Clinic and Research Center, Nepal

²National Zoonoses and Food Hygiene Research Centre, Nepal

³Nepal Academy of Science and Technology, Nepal

⁴Chitwan Medical College and Teaching Hospital, Department of Microbiology, Nepal

⁵Kathmandu College of Science and Technology, Nepal

⁶Department of Science and Technology, Far western University, Nepal

⁷Sukraraj Tropical and Infectious Disease Hospital, Nepal

Article Information

Received date: Oct 03, 2018

Accepted date: Nov 26, 2018

Published date: Nov 27, 2018

*Corresponding author

Yogendra Shah, Everest International Clinic and Research Center, National Zoonoses and Food Hygiene Research Centre, Kathmandu, Nepal,
Tel: +977-9849610127;
Email: yogendra90@hotmail.com

Distributed under Creative Commons CC-BY 4.0

Rabies is a zoonotic viral disease caused by rabies virus, of the genus *Lyssavirus*, family *Rhabdoviridae* [1]. It is one of neglected zoonotic diseases and remains a major public health concern globally. The disease is highly risk in developing countries like Nepal [2-4]. It is an endemic and priority zoonotic disease in Nepal. Rabies is transmitted by bite of infected dogs, cats, monkeys, wild animals (mongoose, wolf, fox, raccoon, jackal and bat) to human population [5,6].

It occurs in two epidemiological cycles: the urban cycle involves domesticated dogs, and the sylvatic cycle involves wild animals [7]. The urban cycle is maintained by the street and community dogs and is the main source of human rabies. The dog population in Kathmandu valley was around 22,300 in 2012 [8]. In countries where dogs are the primary source of infection to humans, vaccination of dogs can help reduce or eliminate the human rabies burden. Vaccines help to establish pre-exposure immunity and to protect individual animals from contracting rabies, hereby preventing further spread to humans or other domestic animals [9].

The first human fatality case due to vampire bat attack was reported from the time of the Spanish colonization of the Americas during the sixteenth century [10]. The previous studies revealed that no human rabies cases transmitted due to bats were documented in Nepal till date [11]. However, a research study showed that endemic colonies of giant fruit bats (*Pteropus giganteus*) known to reservoirs for multiple zoonotic viruses in Kathmandu Valley [12].

Although human cases in developing countries have been mostly associated with dog bites, bat species may also be infected by rabies virus (RABV) [13-14]. Deforestation has drastically reduced the number of natural prey for bats. The migration pattern of bats from rural to urban areas for food supply may also increase potential contact with domestic and wild animal populations and human beings [15]. In some European countries, Cats have been considered a high-risk species for rabies transmission to humans due to predatory behavior of flying birds including bats which may connect rabies from the sylvatic-aerial cycle to urban settings. Such scenarios may appear in major urban areas of Nepal [16].

Rabies is a 100% vaccine preventable disease. Successful elimination of human rabies requires a multisectoral collaborative approach. Prevention of animal rabies, better public awareness and improved access to cost-effective and high-quality human rabies vaccines are essential for the elimination of human rabies [17,18]. However, Nepal has been facing a number of barriers-technical, intersectoral, organizational and financial. In addition, there has been poor implementation sylvatic rabies surveillance, dog rabies control campaigns and dog population control programmes [17,18].

Government of Nepal should take quick and urgent action to prevent the zoonotic rabies virus disease by formulating rabies elimination strategies such as [6]; developing the protocol for identification, characterization of rabies virus at regional, national and international level laboratory, identification and control the source of entry virus, enhancement of laboratory-based surveillance in domestic and wild animals, increased rate of vaccination in animals against rabies, restriction of animal movement, coordination of a multi-agency response [19], provision for public/ professional outreach and education guidelines in addition to implementing the strong strategy plan to further reduce of zoonotic rabies outbreak in future. Globally, one of successful model like One health approach or Ecohealth approach or One Science should be engaged to prevent zoonotic disease by government of Nepal to avoid panic outbreak in future of particular zoonotic rabies virus.

References

1. Knobel DL, Cleaveland S, Coleman PG, Fèvre EM, Meltzer MI, Miranda ME, et al. Re-evaluating the burden of rabies in Africa and Asia. *Bull. World Health Organ.* 2005; 83: 360-368.
2. Karki S, Thakuri KC. Epidemiological situation of animal rabies and its control strategy in Nepal. In: *Proceedings of the 9th Conference of Nepal Veterinary Association*. Kathmandu. 2010; 105-110.
3. Yousaf MZ, Qasim M, Zia S, Khan Mu, Ashfaq UA, Khan S. Rabies molecular virology, diagnosis, prevention and treatment. *Virol J.* 2012; 9: 50.
4. OIE. Animal Health in the World, Rabies Portal. 2018.
5. Condori-Condori RE, Streicker DG, Cabezas-Sanchez C, Velasco-Villa A. Enzootic and epizootic rabies associated with Vampire bats, Peru. *Emerg Infect Dis.* 2013; 19: 1463-1469.
6. Singh R, Singh KP, Cherian S, Saminathan M. Rabies-epidemiology, pathogenesis, public health concerns and advances in diagnosis and control: a comprehensive review, *Veterinary Quarterly.* 2017; 37: 1212-1251.
7. Devleesschauwer B, Aryal A, Sharma BK. Epidemiology, Impact and Control of Rabies in Nepal: A Systematic Review. *PLoS Negl Trop Dis.* 2016; 10: e0004461.
8. Kakati K. Street Dog Population Survey, Kathmandu. In *Final Report to the World Society for the Protection of Animals*. 2012.
9. Berndtsson LT, Nyman AKJ, Rivera E, Klingeborn B. Factors associated with the success of rabies vaccination of dogs in Sweden. *Acta Vet Scand.* 2011; 53: 22.
10. De Oviedo, Valdes F. *Sumario de la Natural historia delas Indias*, 1526. Mexico City: Fond de Cultura Economica. 1950.
11. Pant GR, Lavenir R, Wong FYK, Certoma A, Larrous F. Recent Emergence and Spread of an Arctic-Related Phylogenetic Lineage of Rabies Virus in Nepal. *PLoS Negl Trop Dis.* 2013; 7: e2560.
12. USAID PREDICT. Promoting global health.
13. Sparkes JL, Fleming PJS, Ballard G, Scott-Orr H, Durr S, Ward MP. Canine rabies in Australia: a review of preparedness and research needs. *Zoonoses Public Hlth.* 2014; 62: 237-253.
14. Cordeiro RA, Duarte NFH, Rolim BN, Soares Júnior FA, Franco ICF, Ferrer LL, et al. The importance of wild canids in the epidemiology of rabies in Northeast Brazil: a retrospective study. *Zoonoses Public Health.* 2016; 63: 486-493.
15. Johnson N, Aréchiga-Ceballos N, Aguilar-Setien A. Vampire bat rabies: ecology. *Epidemiology and Control Viruses.* 2014; 6: 1911-1928.
16. Frymus T, Addie D, Belák S, Boucraut-Baralon C, Egberink H, Gruffydd-Jones T, et al. Feline rabies. ABCD guidelines on prevention and management. *J Feline Med Surg.* 2009; 11: 585-593.
17. WHO 2005. The control of neglected zoonotic diseases. In *A route to poverty alleviation*. Geneva: World Health Organisation. 2005; 54.
18. WHO. 2018.
19. Seneschall C, Luna-Farro M. Controlling rabies through a multidisciplinary, public health system in Trujillo, La Libertad, Peru. *Pathog Glob Health.* 2013; 107: 361-366.