

First Record of *Culex (Culex) Coronator* (Diptera: Culicidae) in Havana, Cuba

Magaly Pérez Castillo¹, Raúl Gonzalez Broche², Iris Peraza Cuesta¹, María Elena Mendizábal Alcalá¹, Karelis Chamizo Herrera¹, Roberto E Molina Torriente¹, Maricely Rodríguez Milián¹, Juan Andrés Bisset Lazcano², Maureen Leyva Silva², María del Carmen Marquetti Fernández^{2*}

¹Entomology laboratory, Provincial Hygiene Center, Cuba

²Vector Control Department, Institute of Tropical Medicine Pedro Kourí, Cuba

Article Information

Received date: Jan 08, 2019

Accepted date: Jan 21, 2019

Published date: Jan 22, 2019

*Corresponding author

María del Carmen Marquetti Fernández,
Vector Control Department, Institute
of Tropical Medicine "Pedro Kourí,
Autopista Noviadel Mediodía Km 6 ½,
La Lisa, Havana, Cuba,
Email: marquetti@ipk.sld.cu

Distributed under Creative Commons
CC-BY 4.0

Keywords Arbovirus; *Culex coronator*;
Zoonosis

Abstract

Culex (Culex) coronator Dyar and Knab was recorded in three of the 15 municipalities surveyed in Havana province, Cuba. *Cx. coronator* larvae were found in six different types of breeding sites. At these breeding sites, this mosquito was associated with other species such as *Culex nigripalpus*, *Theobald* and *Anopheles albimanus*, Wiedemann. This represents the first record of *Cx. coronator* in the Havana province.

Introduction

Culex coronator is a neotropical mosquito species described by [1] in Trinidad. Its habitat extended from United State to South America (Mexico, Belize, Guatemala, Honduras, Nicaragua, Costa Rica, El Salvador Panama, Argentina, Colombia, Bolivia, Brazil, French Guiana, Paraguay, Peru, Suriname, Venezuela, and in the Caribbean region in Trinidad and Tobago) [2-3]. In United States, this species is considered an invasive species that had dispersed eastward [2-6]. *Cx. coronator* breeds in a large range of larval habitats that vary from natural to artificial containers, temporary ground pools, under complete and partial shade or full sun, in sylvatic, rural and urban environments [7]. In Brazil, it was found in anthropic environments [8].

The *Cx. coronator* complex has a confuse taxonomy history that includes several species. Recent evidence from the morphological and molecular data concludes, (at least provisionally), that the *Cx. coronator* complex is a single polymorphic species [9].

Cx. coronator has been documented in a number of temperate areas in the United States. Since 2002 specimens have been reported in different geographic areas such as, Oklahoma, Louisiana, Mississippi, Alabama, and Florida. These records extend the known distribution of *Cx. coronator* from six other states (Arizona, Louisiana, Mississippi, New Mexico, and Texas). In 2007, adult specimen's of *Cx. coronator* were collected in Chatham County, Georgia, and Jasper County, South Carolina, during disease vector monitoring efforts [10]. This represents the first Atlantic coast record of this species in Georgia and the first confirmed record of *Cx. coronator* in South Carolina. Subsequently a single adult female of *Cx. coronator* was collected on November 1, 2016, in Suffolk, Virginia, in a BG-Sentinel 2[®] trap during routine mosquito surveillance. This was the first documented observation of this species in the state of Virginia and the first time it has been found this far north in the United States [11]. The distribution of *Cx. coronator* in the United State seems to be expanding at a prodigious rate, for reasons that remain unclear at first. Because *Cx. coronator* was not included in the taxonomic keys of Florida mosquitoes [12], it is possible that it was present in Florida prior to the first report in 2005 [5] and that the mosquito was present in the counties reported here prior to the year that the first confirmed collections and identification of this species were made. Several factors that may have facilitated the spread and establishment of this mosquito in Florida were discussed recently [13].

Although *Cx. coronator* is not usually considered to be a species of major health importance, several pathogens have been isolated from field-collected females. Venezuelan equine encephalitis virus samples from Mexico [14-15] and St. Louis encephalitis virus samples have been isolated from females of *Cx. coronator* in Trinidad [16]. Studies on West Nile virus in United States found this mosquito naturally infected [17] while another study demonstrated its high susceptibility to this virus in laboratory conditions between 80-100% and similar or higher transmission rates than *Culex pipiens*, *Culex quinquefasciatus* and *Culex Restuans* [18].




Figure 1: Municipalities with *Culex coronator* presence in Havana, Cuba, 2014-2018.

In 1981, the eradication campaign of *Aedes aegypti* was established in Cuba, later this campaign received the name of National Control Program of *Ae. aegypti* and *Aedes albopictus*. In both, the identification of mosquito larvae and adults routinely collected is carried out by the personnel belonging to entomology laboratory at municipality and provincial level in whole country. In 1995 was reported *Ae. albopictus* in Cuba by this program [19].

The aim of this paper is to report the presence and distribution of *Cx. coronator* in Havana, Cuba and to identify its breeding sites and association with other mosquito species.

Methodology

Havana is located at 23°07'00"N 82°23'00"W of Cuba. The finding occurred during the sampling was carried out as part of routine mosquito monitoring conducted by the national control program established in the country in each 15 municipalities in the Havana province. The containers and natural breeding sites present in houses, courtyards and surrounding houses were inspected for

the presence of mosquito larvae monthly in all municipalities from January 2014 to June 2018. The work was performed by trained vector control personnel from the national vector control program in each municipality. The larval mosquito samples were collected using a 3mL plastic pipette and placed in vials with 70% alcohol. The vials were labeled with the date, sample collection site, and container type. A key for the mosquito was used to identify the samples [20] in the provincial entomology laboratory and quality control of this process was carried out in the Vector Control Department in the Institute of Tropical Medicine Pedro Kourí.

Results and Discussion

Cx. coronator was found in three (20%) of the 15 surveyed municipalities (the first register was in Habana del Este in 2014 following by Guanabacoa in 2016 and Plaza de la Revolución in 2017) (Figure 1). This mosquito was collected from water storage tanks, cans, cement pots, hole in ground and bucket associated with *An. albimanus* and *Cx. nigripalpus* (Table 1). This represents the first record of *Cx. coronator* in Havana province and in Cuba.

Table 1: General information on *Cx. coronator* reported in three municipalities in Havana province, Cuba, 2014-2018.

Collected Date	Municipality	Locality	Breeding site type	Mosquito species associated
23/07/2014	Habana del Este	Guanabo	cans	4 Larvae of <i>Cx. coronator</i> 2 Larvae of <i>Cx. nigripalpus</i>
2/11/2016	Guanabacoa	Santa María del Rosario	drain	4 Larvae of <i>Cx. coronator</i> 4 Larvae of <i>An. albimanus</i>
15/02/2017	Plaza de la Revolución	Parque Zoológico de La Habana	Cementpots	2 Larvae of <i>Cx. coronator</i> 4 Larvae of <i>An. albimanus</i>
26/06/2018	Guanabacoa	Santa María del Rosario	Hole in ground	6 Larvae of <i>Cx. coronator</i>
26/06/2018	Guanabacoa	Santa María del Rosario	cans	3 Larvae of <i>Cx. coronator</i>
4/7/2018	Guanabacoa	Santa María del Rosario	Water storage tanks	4 Larvae of <i>Cx. coronator</i> 2 Larvae of <i>Cx. nigripalpus</i>
19/07/2018	Guanabacoa	Santa María del Rosario	Bucket	3 Larvae of <i>Cx. coronator</i>
19/07/2018	Guanabacoa	Santa María del Rosario	Water storage tanks	5 Larvae of <i>Cx. coronator</i>
				3 Larvae of <i>Cx. coronator</i>

According to the distribution of this species in the American continent, mainly from the United States to Patagonia, it is difficult to identify the factors that could facilitate the occurrence of *Cx. coronator* in Cuba. The authors believe that the species may have been introduced through adult forms aboard maritime transports (cruises or others) or aboard airplanes from the continental distribution area of the mosquito. Although the possibility remains that *Cx. coronator* was present in the country before and that because of the small amount of taxonomic sampling on the fauna of Cuban culicids, researchers did not detect it until now. In another hand *Cx. coronator* was not included in the taxonomic keys of Cuba mosquitoes.

Study Limitations

The larval samples received at the provincial reference laboratory from other mosquito species other than *Aedes aegypti* and *Aedes albopictus* only account for 10% of what is collected during the routine monitoring of the Program, which influences the number of samples, frequency of appearance and number of larvae of *Cx. coronator* identified until now.

No adult specimens were obtained from the locations where the larval presence of the mosquito was recorded. However, the provincial government program plans to collect more specimens including *Cx. coronator* in the adult phase, as well as, of other associated species in future studies in the mentioned locations in Havana province, Cuba.

Acknowledgments

The authors wish to thank all the vector control staff of National Control Program of *Aedes aegypti* and *Aedes albopictus* in each municipality in Havana province for their contribution to the realization of this work.

References

- Dyar HG, Knab F. The larvae of Culicidae classified as independent organisms. *J NY Entomol Soc.* 1906; 14: 169-230, 242.
- Pecor JE, Harbach RE, Peyton EL, Roberts DR, Rejmankova E, Manguin S, et al. Mosquito studies in Belize, Central America: records, taxonomic notes, and a checklist of species. *J Am Mosq Control Assoc.* 2002; 18: 241-276.
- Bond JG, Casas-Martínez M, Quiroz-Martínez H, Novelo-Gutiérrez R, Marina CF, Ulloa A, et al. Diversity of mosquitoes and the aquatic insects associated with their oviposition sites along the Pacific coast of Mexico. *Parasit Vectors.* 2014; 7: 41.
- Debboun M, Kuhr DD, Rueda LM, Pecor JE. First Record of *Culex* (*Culex*) *coronator* in Louisiana, USA. *J Am Mosq Control Assoc.* 2005; 21: 455-457.
- Smith JP, Walsh JD, Cope EH, Tennant RAJ, Kozak JA III, Darsie RF Jr. *Culex coronator* Dyar and Knab: a new Florida species record. *J Am Mosq Control Assoc.* 2006; 22: 330-332.
- Gray KM, Burkett-Cadena ND, Eubanks MD. Distribution Expansion of *Culex coronator* in Alabama. *J Am Mosq Control Assoc.* 2008; 24: 585-587.
- Moulis RA, Russell JD, Lewandowski HB Jr, Thompson PS, Heusel JL. *Culex coronator* in Coastal Georgia and South Carolina. *J Am Mosq Control Assoc.* 2008; 24: 588-590.
- Ribeiro AF, Urbinatti PR, Duarte AMRC, Paula MB, Pereira DM, Mucci LF, et al. Mosquitoes in degraded and preserved areas of the Atlantic Forest and potential for vector-borne disease risk in the municipality of São Paulo, Brazil. *J Vector Ecol.* 2012; 37: 316-324.
- Laurito M, Briscoe AG, Almiron WR, Harbach RE. Systematics of the *Culex coronator* complex (Diptera: Culicidae): morphological and molecular assessment. *Zoological Journal of the Linnean Society.* 2018; 182: 735-757.
- Kelly R, Mead D, Harrison BA. Discovery of *Culex coronator* Dyar and Knab (Diptera: Culicidae) in Georgia. *Proc Entomol Soc Wash.* 2008; 110: 258-261.
- Akaratovic KI, Kiser JP. First Record of *Culex coronator* in Virginia, with Notes on Its Rapid Dispersal, Trapping Methods, and Biology. *J Am Mosq Control Assoc.* 2017; 33: 225-228.
- Darsie RF Jr, CD Morris. Keys to the adult females and fourth instar larvae of the mosquitoes of Florida (Diptera, Culicidae). *Tech Bull Fl Mosq Contr Assoc.* 2003; 1: 1-159.
- Connelly CR, Alto BW, O'Meara GF. The spread of *Culex coronator* (Diptera: Culicidae) throughout Florida. *J Vector Ecol.* 2016; 41: 195-199.
- Scherer WF, Dickerman RW, Diaz-Najera A, Ward BA, Miller MH, Schaffer PA. Ecologic studies of Venezuelan encephalitis virus in southeastern Mexico. 3. Infection of mosquitoes. *Am J Trop Med Hyg.* 1971; 20: 969-979.
- Burguete J, Romero-Acevedo S, Salido RF, Pierce EP. Epizootic and epidemic of Venezuelan equine encephalitis in the state of Morelos. *Salud Pub de Mex.* 1973; 15: 231-235.
- Anderson CR, Aitken TH, Downs WG, Spence L. The isolation of St. Louis virus from Trinidad mosquitoes. *Am J Trop Med Hyg.* 1957; 6: 688-692.
- Mackay AJ. Detection of West Nile virus activity in male and female mosquitoes, and evaluation of host utilization patterns of mosquitoes, in East Baton Rouge Parish, Louisiana. Louisiana State University, Baton Rouge, LA, United States in 2007.
- Alto BW, Connelly CR, O'Meara GF, Hickman D, Karr N. Reproductive biology and susceptibility of Florida *Culex coronator* to infection with West Nile Virus. *Vector Borne Zoonotic Dis.* 2014; 14: 606-614.
- Broche RG, Borja EM. *Aedes albopictus* in Cuba. *J Am Mosq Control Assoc.* 1999; 15: 569-570.
- Zapata-Peniche A, Manrique-Saide PC, Rebollar-Téllez EA, Che-Mendoza A, Dzúl-Manzanilla F. Identificación de larvas de mosquitos (Diptera: Culicidae) de Mérida, Yucatán, México y sus principales criaderos. *Rev Biomed.* 2007; 18: 3-17.