

Unravelling Rhizospheric Diversity and
Potential of Phytase Producing MicrobesVinod Kumar*, Ajar Nath Yadav, Abhishake Saxena, Punesh Sangwan and
Harcharan Singh Dhaliwal

Department of Biotechnology, Akal College of Agriculture, Eternal University, Sirmour-173101, India

Article Information

Received date: Mar 11, 2016

Accepted date: Mar 12, 2016

Published date: Mar 15, 2016

*Corresponding author

Vinod Kumar, Department of
Biotechnology, Akal College of
Agriculture, Eternal University,
Sirmour-173101, India, Email: sangwan.
vinod@yahoo.com

Distributed under Creative Commons
CC-BY 4.0

Phosphorus (P) is considered the ultimate limiting nutrient for plants because of its form as insoluble complexes. To address P deficiency, different organic and inorganic fertilizers are added to soils. Inorganic P instantly become unavailable by forming complexes with metal ions and excess P-fertilizers application also leads to water eutrophication of P. Phytic Acid (PA; constitutes 15-50% of total P depending on soil types) as a component of organic P also form stable complexes and its natural degradation is almost impossible. Therefore, role of P solubilizing microbes in rhizosphere become important for P cycling. Several rhizospheric bacterial and fungal species have been reported to play important role in P solubilization in soil [1]. Based on several studies on diversity of P solubilizing microbes and enzymes in varying agro-ecosystems, soil P availability and its uptake by plants is mainly attributed to microbial phosphatases and plant exudates. Phytase producing microbes have gained recent interest due to their plant growth promoting abilities and P pollution management applications [2,3].

Several research groups have studied diversity of different groups of phytases and organisms in diverse habitats for multiple applications of phytases and it includes phytase producing microbes in rumen by Yanke, et al. [2] and Nakashima, et al. [4], β -propeller phytases in aquatic environment by Cheng and Lim [5], phytase producing marine yeasts by Hirimuthugoda, et al. [6], phytase producing yeasts in fermented food by Hellström, et al. [7], and β -propeller phytase genes in the intestinal contents of grass carps by Huang, et al. [8]. A detailed bioinformatic analysis for diversity of phytases was carried out by Lim, et al. [9], and the comparative 'in silico' gene analysis of phytases was carried out by Kumar, et al. [10-12]. Phytase producing bacterial isolates were also reported from rhizosphere of pasture plants. Jorquera, et al. [13] and Acuña, et al. [14] suggested more than one mechanism for utilizing insoluble forms of P. After a long term experiment to study the effect of organic and conventional cropping systems on crop performance and soil fertility, Mäder, et al. [15], suggested that the higher soil microbiological activity and higher biodiversity found in organic plots. To understand the complexity of P availability and uptake, Oberson, et al. [16] have studied fresh and residual phosphorus uptake by ryegrass from soils with different fertilization histories. Some others were also carried out to study effect of varying fertilization and soil conditions on microbial diversity [1,17-19]. Considering the fact that plants are not able to utilize P directly from organic P, studies has been conducted with the objectives of determining the effect of phytase producing bacteria on plant growth [20-22]. Several phytase producing bacterial isolates from Himalayan soils were reported with plant growth promoting activities [1,23]. Rangarajan, et al. [24] has demonstrated change in diversity of P solubilizing pseudomonads isolated from three different plant rhizospheres and concluded variation in soil was more predominant on changing diversity than plant types. Azziz, et al. [25] have studied abundance and diversity of culturable phosphate solubilizing bacteria (PSB) under crop rotations and found potential isolates for use as bioinoculants. Effect of long-term application of manure and fertilizer on biological and biochemical activities (including phosphatase activity) in soil during crop development stages was studied by Mandal, et al. [26], and an interaction between grain yield and phosphatase activity was reported. Genetic diversity of plant growth promoting rhizobacteria isolated from rhizospheric soil of wheat under saline condition was studied by Verma, et al. [27]. According to a study by Nayak, et al. [28], long-term application of compost influences microbial biomass and enzyme activities in a tropical *Aeric Endoaquept* planted to rice under flooded condition. Role of phytase in PA rich soils have not been explored and a study for diversity of these microbes (culturable and un-culturable) and phytase activity in different agro-ecosystems might reveal importance of this unique community in understanding of underlying plant-microbe interactions.

References

1. Kumar V, Singh P, Jorquera MA, Sangwan P, Kumar P, Verma A, et al. Isolation of phytase-producing bacteria from Himalayan soils and their effect on growth and phosphorus uptake of Indian mustard (*Brassica juncea*). World J Microbiol Biotechnol. 2013; 29: 1361-1369.

2. Kumar V, Singh D, Sangwan P, Gill PK. Management of Environmental Phosphorus Pollution Using Phytases: Current Challenges and Future Prospects. *Applied Environmental Biotechnology: Present Scenario and Future Trends*. Springer. 2015; 97-114.
3. Yanke L, Bae H, Selinger L, Cheng K. Phytase activity of anaerobic ruminal bacteria. *Microbiology*. 1998; 144: 1565-1573.
4. Nakashima BA, McAllister TA, Sharma R, Selinger LB. Diversity of phytases in the rumen. *Microbial ecology*. 2007; 53: 82-88.
5. Cheng C, Lim BL. Beta-propeller phytases in the aquatic environment. *Archives of microbiology*. 2006; 185: 1-13.
6. Hirimuthugoda NY, Chi Z, Li X, Wang L, Wu L. Diversity of phytase-producing marine yeasts. *Ciencias Marinas*. 2006; 32: 673-682.
7. Hellström AM, Vázquez-Juárez R, Svanberg U, Andlid TA. Biodiversity and phytase capacity of yeasts isolated from Tanzanian togwa. *International journal of food microbiology*. 2010; 136: 352-358.
8. Huang H, Shao N, Wang Y, Luo H, Yang P, Zhou Z, et al. A novel beta-propeller phytase from *Pedobacter nyackensis* MJ11 CGMCC 2503 with potential as an aquatic feed additive. *Applied microbiology and biotechnology*. 2009; 83: 249-259.
9. Lim BL, Yeung P, Cheng C, Hill JE. Distribution and diversity of phytase-mineralizing bacteria. *The ISME journal*. 2007; 1: 321-330.
10. Kumar V, Singh G, Verma AK, Agrawal S. *In Silico* characterization of histidine acid phytase sequences. *Enzyme research*. 2012; 2012: 845465.
11. Kumar V, Singh G, Sangwan P, Verma AK, Agrawal S. Cloning, Sequencing, and *In Silico* Analysis of Propeller Phytase *Bacillus licheniformis* Strain PB-13. *Biotechnology research international*. 2014; 2014: 1-11.
12. Kumar V, Agrawal S. An insight into protein sequences of PTP-like cysteine phytases. *Nusantara Bioscience*. 2014; 6: 102-106.
13. Jorquera MA, Hernández MT, Rengel Z, Marschner P, de la Luz Mora M. Isolation of culturable phosphobacteria with both phytate-mineralization and phosphate-solubilization activity from the rhizosphere of plants grown in a volcanic soil. *Biology and Fertility of Soils*. 2008; 44: 1025-1034.
14. Acuña J, Jorquera M, Martínez O, Menezes-Blackburn D, Fernández M, Marschner P, et al. Indole acetic acid and phytase activity produced by rhizosphere bacilli as affected by pH and metals. *Journal of soil science and plant nutrition*. 2011; 11: 1-12.
15. Mäder P, Fliessbach A, Dubois D, Gunst L, Fried P, Niggli U. Soil fertility and biodiversity in organic farming. *Science*. 2002; 296: 1694-1697.
16. Oberson A, Tagmann HU, Langmeier M, Dubois D, Mäder P, Frossard E. Fresh and residual phosphorus uptake by ryegrass from soils with different fertilization histories. *Plant and Soil*. 2010; 334: 391-407.
17. Marschner P, Crowley D, Yang CH. Development of specific rhizosphere bacterial communities in relation to plant species, nutrition and soil type. *Plant and Soil*. 2004; 261: 199-208.
18. Yuan W, Zhao X, Ge X, Bai F. Ethanol fermentation with *Kluyveromyces marxianus* from Jerusalem artichoke grown in salina and irrigated with a mixture of seawater and freshwater. *Journal of applied microbiology*. 2008; 105: 2076-2083.
19. He J-S, Wang L, Flynn DF, Wang X, Ma W, Fang J. Leaf nitrogen: phosphorus stoichiometry across Chinese grassland biomes. *Oecologia*. 2008; 155: 301-310.
20. Yadav R, Tarafdar J. Phytase and phosphatase producing fungi in arid and semi-arid soils and their efficiency in hydrolyzing different organic P compounds. *Soil Biology and Biochemistry*. 2003; 35: 745-751.
21. Hariprasad P, Niranjana S. Isolation and characterization of phosphate solubilizing rhizobacteria to improve plant health of tomato. *Plant and Soil*. 2009; 316: 13-24.
22. Singh B, Satyanarayana T. Microbial phytases in phosphorus acquisition and plant growth promotion. *Physiology and Molecular Biology of Plants*. 2011; 17: 93-103.
23. Singh P, Kumar V, Agrawal S. Evaluation of phytase producing bacteria for their plant growth promoting activities. *International Journal of Microbiology*. 2014; 2014: 426483.
24. Rangarajan S, Loganathan P, Saleena LM, Nair S. Diversity of pseudomonads isolated from three different plant rhizospheres. *Journal of applied microbiology*. 2001; 91: 742-749.
25. Azziz G, Bajsa N, Haghjou T, Taulé C, Valverde Á, Igual JM, et al. Abundance, diversity and prospecting of culturable phosphate solubilizing bacteria on soils under crop-pasture rotations in a no-tillage regime in Uruguay. *Applied soil ecology*. 2012; 61: 320-326.
26. Mandal A, Patra AK, Singh D, Swarup A, Mastro RE. Effect of long-term application of manure and fertilizer on biological and biochemical activities in soil during crop development stages. *Bioresource Technology*. 2007; 98: 3585-3592.
27. Verma P, Yadav AN, Kazy SK, Saxena AK, Suman A. Molecular diversity and multifarious plant growth promoting attributes of Bacilli associated with wheat (*Triticum aestivum* L.) rhizosphere from six diverse agro-ecological zones of India. *J Basic Microbiol*. 2016; 56: 44-58.
28. Nayak DR, Babu YJ, Adhya T. Long-term application of compost influences microbial biomass and enzyme activities in a tropical *Aeric Endoaquept* planted to rice under flooded condition. *Soil Biology and Biochemistry*. 2007; 39: 1897-1906.