

A Case of Koolen de vries Syndrome or 17q21.31 Microdeletion Syndrome Associated with Infertility: A Case Report

Hojatolah Rezaei¹, Neda Asgharzadeh² and Azra Moradbeigi³

¹Devision for counselling genetics, Babol Medical University, Iran,

²Shomal Infertility Clinic, Amol, Iran

³Azra Moradbeigi, Counselor in Welfare Organization of Qom, Iran

Article Information

Received date: Feb 02, 2018

Accepted date: April 16, 2018

Published date: Apr 30, 2018

*Corresponding author

Hojatolah Rezaei, Devision for counselling genetics, Babol Medical University, Babol, Iran,

Tel: 09127578537-00981132289084;

Email: Hojatrezaei60@gmail.com

Distributed under Creative Commons CC-BY 4.0

Keywords Koolen de Vries or 17q21.31 microdeletion syndrome; Array-CGH; Infertility

Abstract

The chromosome 17q21.31 microdeletion syndrome (Koolen de vries Syndrome) is a genomic disorder in patients with unexplained mental retardation that has originally been identified using high resolution genome analyses such as array CGH. Here we report the clinical characterization of a man with the 17q21.31 microdeletion syndrome and infertility. Clinical examination reveals that mild developmental delay, poor speech development, stutter, facial dysmorphisms including a long face, high narrow palate, a tubular or pear-shaped nose and a bulbous nasal tip, a friendly behaviour and the main characteristic features to refer was infertility. Other clinically important features include hypothyroidism, bilateral asymmetry in brain MRI, without any anomalies in urogenital examination and echocardiography. Using array CGH we found pathologic loss of 921 kb on 17q21.31 compatible with Koolen de Vries syndrome. The chromosome 17q21.31 microdeletion syndrome or Koolen de Vries syndrome recognizable as a genomic disorder with broad features and infertility may be occur.

Introduction

New technologies such as Array Comparative Genomic Hybridization (aCGH), have paved the way for identifying, a small but significant percentage of genomic disorders before their clinical description [1]. One of these cases is the chromosome 17q21.31 microdeletion syndrome (Mendelian Inheritance in Man (MIM) 610443) or Koolen de Vries Syndrome (KDVS) characterized by a recurrent 500-650 kb deletion [2]. Screening of large heterogeneous cohorts of individuals with Mental Retardation (MR) using aCGH led to identifying the recurrent microdeletion for first time [2]. The prevalence of this syndrome was estimated to be 1 in 16,000 individuals and most important features including developmental delay, a friendly/amiable behavior, hypotonia, and facial dysmorphisms such as a long face, a tubular or a 'pear-shaped' nose with a bulbous nasal tip, epilepsy, heart defects and urologic abnormalities [3]. The critical region that is missing in most individuals with KDVS is 424kb encompasses at least six genes and KANSL1 gene has been shown to make a major contribution to the phenotype [3].

Here, we report on an MR patient referred to genetics clinic by an infertility & reproductive center.

Case report

A 28 year old man with mental delay development referred to genetics clinic by the infertility & reproductive center. He was initially referred to an infertility & reproductive center as he was infertile. The proband was the last born boy to non-consanguineous parents and without any history of MR and infertility in the family. He was born in good condition and did not require resuscitation without any prenatal/postnatal problems such as neonatal hypoxia and seizure. At the age of two years he was further referred to a pediatric clinic because of poor Verbal skills. It was noted that he still had poor speech development. Developmental delay was further confirmed some years later. The patient has a friendly behavior and mild intellectual disability. Clinical examination revealed sparse thick hair, a triangular face with a high flat forehead, ptosis, a bulbous nasal tip, high narrow palate, widely space nipple, scoliosis and ectodermal abnormalities such as multiple nevi, depigmentosa, hemangiomas. Her fingers are slender and long with hypoplastic muscles. In addition bilateral asymmetry was shown by brain MRI. Semen analysis or seminogram have been shown Oligozoospermia (Very low sperm count), Asthenozoospermia (poor sperm motility), Teratozoospermia (sperm carry more morphological defects than usual). He has hypothyroidism, but levels of the prolactin and testosterone hormones is normal.

Cytogenetic and Molecular Studies

Cytogenetic karyotype of peripheral blood lymphocytes by GTG-banding, at the 450-550 showed a normal 46, XY karyotype. However molecular analysis, array-CGH, showed the presence of chromosomal alteration. Whole genome oligo array CGH was performed using CYTOCHIP ISCA 8X60K whole genome oligo array version 2 and was analyzed using Blue Fuse Multi Software. The array consists of 60000 spots with average backbone resolution of 51 Kbs and close to 500 targeted disease regions. The sample was hybridized twice against normal samples as references. Array-CGH analysis identified an interstitial deletion of a segment of ~921 kb on 17q21.31q21.31 from nucleotide 43,706,915 to 44,628,121 compatible with 17q21.31 microdeletion syndrome KDVS which is a contiguous gene syndrome caused by microdeletion (600-800 kb) of chromosome 17q21.31 encompassing CRHR1(122561), MAPT(157140), STH (607067), IMP5 (608284) and KANSL1(612452).

Discussion

Mental retardation occurs in approximately 2-3 % of the general population [4].

The 17q21.31 microdeletion syndrome is a new delay developmental syndrome [2,5]. Only a few cases can be detected based on the clinical features of the patient, so individuals with 17q21.31 microdeletion syndrome is currently highly underdiagnosed and in majority of cases the microdeletion was identified after array-CGH [3]. Although microdeletion size can vary, most of them are between 500-650 kb [5]. A minimal chromosomal region has been refined to a segment of 424kb at chr17:41046729-41470954 (hg 17) and includes the MAPT, CRHR1, KANSL1, SPPL2C, and STH genes [3]. A number of recent reports have shown that haploinsufficiency of KANSL1 gene is sufficient to cause the full KDVS phenotype [6,7]. All of the signs and symptoms our patient presents have previously been seen in reported cases of the 17q21.31 microdeletion syndrome, with the exception of infertility. There is no family history of infertility and ultrasounds indicate that the genitourinary system is healthy in our patient. Here, we reported a patient with one of the largest microdeletions ever described which consists of approximately 921 kb and contains the following genes: MAPT, CRHR1, KANSL1, SPPL2C and STH, the hypothetical ORF LOC644246 and the non-coding RNA MGC57346.

So far, for variety of reasons, there is no report that individual with KDVS is known to have had children of their own. For example because the lack of identification of affected people so remain undiagnosed or less likely to have children than normal people.

Haploinsufficiency of one or more other elements within the critical region or position effect resulting 17q21.31 with effect on gametogenesis perhaps representing the infertility seen in the KDVS syndrome. Because structural complexity of this genomic region and due to unknown the exact function of this region elements, infertility traits in individuals with KDVS can be challenging.

Conclusion

Here we reported a patient with KDVS associated infertility. No firm conclusions can be made about infertility because further clinical studies and more DNA testing in other patients with KDVS are needed.

References

1. Slavotinek AM. Novel microdeletion syndromes detected by chromosome microarrays. *Hum. Genet.* 2008; 124: 1-17.
2. Koolen DA, Vissers LE, Pfundt R, de Leeuw N, Knight SJ, Regan R, et al. A new chromosome 17q21.31 microdeletion syndrome associated with a common inversion polymorphism. *Nat Genet.* 2006; 38: 999-1001.
3. Koolen DA, Sharp AJ, Hurst JA, Firth HV, Knight SJL, Goldenberg A, et al. Clinical and molecular delineation of the 17q21.31 microdeletion syndrome. *J Med Genet.* 2008; 45: 710-720.
4. Leonard H, Wen X. The epidemiology of mental retardation: Challenges and opportunities in the new millennium. *Ment Retard Dev Disabil Res Rev.* 2002; 8:117-134.
5. Sharp AJ, Hansen S, Selzer RR, Cheng Z, Regan R, Hurst JA, et al. Discovery of previously unidentified genomic disorders from the duplication architecture of the human genome. *Nat Genet.* 2006; 38:1038-1042.
6. Koolen DA, Kramer JM, Neveling K, Nillesen WM, Moore-Barton HL, Frances VE, et al. Mutations in the chromatin modifier gene KANSL1 cause the 17q21.31 microdeletion syndrome. *Nat Genet.* 2012; 44: 639-641.
7. Zollino M, Orteschi D, Murdolo M, Lattante S, Battaglia D, Stefanini C, et al. Mutations in KANSL1 cause the 17q21.31 microdeletion syndrome phenotype. *Nat Genet.* 2012; 44: 636-638.
8. Koolen DA, Dupont J, de Leeuw N, Lisenka ELMV, van den Heuvel SPA, Alyson B, et al. Two families with sibling recurrence of the 17q21.31 microdeletion syndrome due to low-grade mosaicism. *Eur J Hum Genet.* 2012; 20: 729-733.