

In-vitro Antimalarial and Antileishmanial Studies of *Markhamia platycalyx* Sprague Leaves

Basma Khalaf Mahmoud¹, Ashraf Nageeb El-Sayed Hamed^{1*}, Mamdouh Nabil Samy¹, Amira Samir Wanas^{1,2}, Mohamed M Radwan², Mahmoud A ElSohly^{2,3}, Mohamed Salah Kamel¹

¹Department of Pharmacognosy, Faculty of Pharmacy, Minia University, 61519 Minia, Egypt

²National Center for Natural Products Research, School of Pharmacy, The University of Mississippi, University, MS 38677, USA

³Department of Pharmaceutics, School of Pharmacy, The University of Mississippi, University, MS 38677, USA

Article Information

Received date: Jul 31, 2017

Accepted date: Aug 26, 2017

Published date: Sep 01, 2017

*Corresponding author

Ashraf Nageeb El-Sayed Hamed,
Department of Pharmacognosy, Faculty
of Pharmacy, Minia University, 61519
Minia, Egypt, Tel: +2086-234-77-59;
Fax: +2086-236-90-75;
Email: ashrafnag@mu.edu.eg

Distributed under Creative Commons
CC-BY 4.0

Keywords *Markhamia platycalyx*;
Bignoniaceae; *In-vitro*; Antimalarial;
Antileishmanial

Article DOI 10.36876/smjmps.1002

Abstract

Bignoniaceae is rich in active metabolites and includes numerous genera of high economic and therapeutic values. One of these plants is *Markhamia platycalyx*. The petroleum ether fraction of the total ethanol leaf extract of *M. platycalyx* exhibited IC₅₀ 26760 ng/mL against *Plasmodium falciparum* D₆ and 22430 ng/mL against *P. falciparum* W₂ in the 2nd phase assay. It did not show any cytotoxic activity against the VERO mammalian cells, indicating the safety of the petroleum ether fraction. Additionally, the dichloromethane fraction of the same extract was the most active fraction against *Leishmania donovani* amastigotes in THP1 with 86% inhibition in the 1st phase assay, which was higher than Amphotericin B.

Introduction

Natural products have extensive significant attention in recent years due to their various pharmacological activities as cytotoxic [1], hepatoprotective [2], anti-inflammatory [3], anti-pyretic [3], gastroprotective [3], anti-diabetic [3],....etc.

The parasitic diseases (malaria, trypanosomiasis, leishmaniasis, schistosomiasis, lymphatic filariasis and onchocerciasis) in tropical regions have been a dangerous public health problem especially in middle- and low-income countries. These diseases affect millions of people, resulting in thousands of death annually [4-5].

Family Bignoniaceae is rich in the secondary metabolites and includes numerous genera of high economic and therapeutic values [6,7]. It is found in tropical and subtropical areas with a few species in temperate climates [8]. It is comprised of 104 genera and 860 species [8]. One of these species is *Markhamia platycalyx* Sprague (Syn. *Dolichandrone platycalyx* Baker). *M. platycalyx* tree, known in Uganda under the native name Lusambia, is said to yield "the finest of local timbers" (Mahon) [9].

By reviewing the literature, some researchers considered *M. platycalyx* as a synonym of *M. lutea*, however, a recent study on cultivated plants in Egypt, classified both of them in two different lineages [10].

By reviewing the antimalarial and antileishmanial literature of the genus; *M. lutea* was investigated for many therapeutic activities such as antimalarial [11], antileishmanial [11] and antiviral [12]. Moreover, the stem bark of *M. tomentosa* had a good antimalarial activity *in-vitro* [13]. While, nothing could be found for *M. platycalyx*. This provoked us to carry out extensive studies on this plant including *viz.*, *in-vitro* antimalarial in addition to antileishmanial activities.

Materials and Methods

Materials

Amphotericin B, Chloroquine, Artemisin and Amikacin (Sigma-Aldrich, USA). Solvents used in this work, e.g. light petroleum ether, chloroform, ethyl acetate, ethanol, methanol and distilled water were purchased from El-Nasr Company for Pharmaceuticals and Chemicals, Egypt.

Apparatus

Rotary evaporator (Laborota Heidolph, Germany) was used for distillation the solvents. Circulating hot-air oven (Carbolite, Germany) was used for drying. Sonicator (Wise Clean WUC

OPEN ACCESS

ISSN: 2638-4698

Table 1: *In-vitro* antimalarial activity (1st screening) of *M. platycalyx* leaves against *P. falciparum* D₆ strain.

Extract/Fraction/Compound	% of Inhibition
TEE	8
Petroleum ether fraction	87
DCM fraction	47
EtOAc fraction	24
Aqueous fraction	14
CP	1
Chloroquine	100
All (15867 ng/mL) except Chloroquine (79 ng/mL)	

A-10H, Dahan Scientific Co. Ltd, Korea) was used for dissolving the materials used. Water distillater (Bhanu Basic/PH4, MK-I, India).

Plant material

The leaves of *M. platycalyx* Sprague were collected in May 2012 from El-Zohria Botanical Garden, Giza, Egypt. The plant was identified by Dr. Mamdouh Shokry, Director of El-Zohria Botanical Garden, Giza, Egypt. A voucher specimen (Mn-Ph-Cog-015) was kept in the Herbarium of Pharmacognosy Department, Faculty of Pharmacy, Minia University, Minia, Egypt.

Preliminary Phytochemical Screening

Preliminary qualitative phytochemical screening of the total ethanol extract (TEE) of *M. platycalyx* leaves was carried out using standard reported phytochemical procedures [14,15].

Extraction and Fractionation of *M. platycalyx* leaves

The air dried powdered leaves (5 kg) of *M. platycalyx* were extracted by maceration with 95% ethanol (3x, 5 L each) and then concentrated under reduced pressure to give (500 g) of a viscous residue. It was suspended in the least amount of distilled water (600 mL), transferred to a separating funnel and partitioned successively with light petroleum ether, dichloromethane (DCM) and ethyl acetate (EtOAc) (each solvent; 3x, 350 mL each). The obtained fractions were concentrated under reduced pressure using rotary evaporator to afford three fractions: petroleum ether (95 g), DCM (19 g), EtOAc (30 g). The remaining aqueous layer was concentrated to afford (250 g).

Preparation of Crude Polysaccharide (CP)

A part of the concentrated aqueous fraction (25 g) was gradually added to a flask containing one liter methanol with vigorous shaking and kept in a refrigerator overnight, then filtrated using Büchner funnel under reduced pressure. The residue was dried in the oven 40°C to yield (3 g) of crude polysaccharide powder.

Table 2: *In-vitro* antimalarial activity (2nd assay) of the petroleum ether fraction of *M. platycalyx* leaves.

Fraction/Compound	<i>P. falciparum</i> D ₆		<i>P. falciparum</i> W ₂		VERO cells
	IC ₅₀	SI	IC ₅₀	SI	IC ₅₀
Petroleum ether fraction (47600- 5289 ng/mL)	26760	> 1.8	22430	> 2.1	> 47600
Artemisin (238-26.4 ng/mL)	< 26.4	> 9	172.6	> 1.4	> 238
Chloroquine (238-26.4 ng/mL)	< 26.4	> 9	< 26.4	> 9	> 238

In-vitro antimalarial activity

Antimalarial activity was estimated in-vitro against chloroquine sensitive (D₆, Sierra Leone) and resistant (W₂, Indo China) strains of *Plasmodium falciparum* by determining Plasmodial Lactate Dehydrogenase (PLDH) activity [16]. A 200 µL suspension of *P. falciparum* culture (2% parasitemia and 2% hematocrit in RPMI 1640 medium supplemented with 10% human serum and 60 µg/mL Amikacin) was added to the wells of a 96-well plate containing 10 µL of serially diluted samples. The plate was flushed with a gas mixture of 90% N₂, 5% O₂ and 5% CO₂ and incubated at 37°C for 72 h in a modular incubation chamber. PLDH activity was determined by using Malstat™ reagent (Flow Inc., Portland, OR). Briefly, 20 µL of the incubation mixture was mixed with 100 µL of the Malstat™ reagent and incubated for 30 min. Then, 20 µL of a 1:1 mixture of NBT/PES (Sigma, St. Louis, MO) was added and the plate is further incubated for an hour in dark. The reaction was stopped by adding 100 µL of a 5% acetic acid solution. The plate was measured at 650 nm using the EL-340 Biokinetics Reader (Bio-Tek Instruments, Vermont). IC₅₀ values were obtained from the dose-response curves generated by plotting percent growth versus drug concentration. Chloroquine was included in each assay as positive control. DMSO (0.25%) was used as a vehicle control. The TEE and different fractions of *M. platycalyx* leaves were initially tested against *P. falciparum* D₆ strain in a 1st phase screening at 15867 ng/mL and the percentage of inhibition was calculated relative to the negative and positive controls. The tested samples that resulted in % inhibition ≥ 50% proceeded to 2nd assay.

In the 2nd phase assay, the tested extract and fractions passing 1st phase screening dissolved to 20 mg/mL and tested at three concentrations 47600, 15867, 5289 ng/mL and IC₅₀ were determined. The Selectivity Index (SI) was calculated. All IC₅₀ were calculated using the XLfit curve. Artemisin and Chloroquine (standard antimalarial drugs) were used as positive controls. All experiments were carried out in duplicate. The results of the antimalarial activity were listed in tables 1 and 2. The *in-vitro* cytotoxicity was also determined against mammalian kidney fibroblasts (VERO cells). The assay was performed in 96-well tissue culture-treated plates as described earlier [17]. Briefly, cells were seeded in the wells of a 96-well plate (25,000 cells/well) and incubated for 24 h. Samples were added and plates were again incubated for 48 h. The number of viable cells was determined by neutral red assay. IC50 values were determined from dose-response curves. Amphoterecin B was used as a positive control, while DMSO was used as vehicle control. The results were showed in table 2.

In-vitro antileishmanial activity

The antileishmanial activity of the TEE and different fractions of *M. platycalyx* leaves were tested *in-vitro* against a culture of *Leishmania donovani* promastigotes [18]. The promastigotes were grown in RPMI 1640 medium supplemented with 10% fetal calf

Table 3: *In-vitro* antileishmaial activity (1st phase screening) of *M. platycalyx* leaves.

Extract/Fraction/Compound	% of Inhibition		
	<i>L. donovani</i> _Pinh	<i>L. donovani</i> AMAST_Pinh	<i>L. donovani</i> AMASTTHP_Pinh
TEE	0	9	13
Petroleum ether fraction	6	8	31
DCM fraction	18	48	86
EtOAc fraction	1	12	0
Aqueous fraction	2	12	0
CP	2	13	5
Amphotericin B	99	96	81
All (20 µg/mL) except Amphotericin B (0.4 µg/mL)			

serum (Gibco Chem. Co.) at 26°C. A 3-day-old culture was diluted to 5×10^5 promastigotes/mL. Drug dilutions were prepared directly in cell suspension in 96-well plates. Plates were incubated at 26°C for 48 h and growth of *L. donovani* promastigotes was determined by the Alamar blue assay. Standard fluorescence was measured on a Fluostar Galaxy plate reader (BMG Lab Technologies) at an excitation wavelength of 544 nm and an emission wavelength of 590 nm. Amphotericin B was used as the standard antileishmanial agents. IC₅₀ values were computed from dose-response curves.

The tested samples that showed % inhibition $\geq 50\%$ proceeded to 2nd assay. In the 2nd assay, the tested extract and fractions were dissolved to 20 mg/mL and tested at (40.0, 8.0 & 1.6 µg/mL). All IC₅₀ and IC₉₀ were calculated using the XLfit curve. The Amphotericin B (standard drug) was used as positive control [19,20]. All experiments were carried out in duplicate. The results of the antileishmanial (1st phase) were listed in Table 3.

Results and Discussion

Preliminary phytochemical screening

The TEE of *M. platycalyx* leaves showed the presence of carbohydrates and/or glycosides, flavonoids, unsaturated sterols and/or triterpenes and anthraquinones. On the other hand, it was free from crystalline sublimable substances, saponins, alkaloids, tannins, cardenolides and coumarins. This preliminary phytochemical screening showed many classes of secondary metabolites indicating various expected biological activities.

In-vitro antimalarial activity

The petroleum ether fraction showed the highest percentage of inhibition (87%) against *P. falciparum* D₆ strain relative to chloroquine. Any drug caused $\geq 50\%$ inhibition; it proceeded to 2nd phase assay. On the other hand, the DCM fraction exhibited a good antimalarial activity 47% inhibition, followed by EtOAc fraction 24%, while the aqueous fraction showed weak activity 14%, but higher than that of the TEE (8%) as shown in Table 1.

The 2nd phase assay of the petroleum ether fraction (Table 2) revealed IC₅₀ of 26760 ng/mL against *P. falciparum* D₆ and 22430 ng/mL against *P. falciparum* W₂. It did not show any cytotoxic activity against the VERO mammalian cells line up to the maximum dose tested; 47600 ng/mL, as recorded in Table 2, indicating the safety of the petroleum ether fraction.

The previous phytochemical review showed various classes of compounds isolated and identified from genus *Markhamia* viz., phenylpropanoids, lignans, naphthoquinones, anthraquinones, sterols, cycloartane triterpenes and their glycoside derivatives, phenolic glycosides and triterpene acids. These compounds are isolated from different plant parts including roots, leaves, stem, root bark and heart wood [21].

Moreover, sterols and triterpenes were found in the preliminary phytochemical screening of TEE. Consequently, petroleum ether has an ability to extract sterols and triterpenes from TEE. Therefore, the activity may be due to presence of sterols and triterpenes in this fraction [22,23]. Our findings are in line with the previous studies [11,13].

In-vitro antileishmanial activity

The 1st phase screening of antileishmanial activity in Table 3 revealed that the DCM fraction was the most active against *L. donovani* amastigotes in THP1 with 86% inhibition, which is even higher than Amphotericin B. While, the other fractions showed weak or no activity. The DCM fraction in 2nd phase assay showed IC₅₀ and IC₉₀ against *L. donovani* amastigote > 20 µg/mL.

Leishmaniasis is a vector-borne disease, affecting 72 developing countries and 13 of the least developed countries. Visceral leishmaniasis due to *L. donovani* is the most severe form of Leishmania infections. The annual incidence of visceral leishmaniasis is estimated to be 500,000 cases. The overall prevalence of visceral leishmaniasis is 12 million people and the population at risk is 350 million [24]. The drug of choice for the treatment is still a problem. Therefore, there is an urgent need to discover new drugs with high activity and low side effects. Natural products have become a key source of new drugs in the last years [19,25].

As mentioned before, the earlier phytochemical review exhibited the presence of different classes of compounds isolated and identified from *Markhamia* species as phenyl propanoids, lignans, naphthoquinones, anthraquinones, sterols, cycloartane triterpenes and their glycoside derivatives, phenolic glycosides and triterpene acids [21]. Moreover, sterols and triterpenes were found in the preliminary phytochemical screening of TEE. Consequently, DCM has an ability to extract sterols and triterpenes from TEE. Therefore, the activity may be due to presence of sterols and triterpenes [20,22]. Our results are in line with the previous studies on another two species [11,13].

Conclusion

In this study, the significant antimalarial and antileishmanial activities make *Markhamia platycalyx* leaves a potential source for the antiprotozoal drugs that are strongly recommended for further development.

References

1. Hamed ANE, Wätjen W, Schmitz R, Chovolou Y, Edrada-Ebel R, Youssef DTA, et al. A New Bioactive Sesquiterpenoid Quinone from the Mediterranean Sea Marine Sponge *Dysidea avara*. *Nat Prod Commun*. 2013; 8: 289-292.
2. Hamed ANE, Wahid A. Hepatoprotective activity of *Borago officinalis* extract against CCl₄-induced hepatotoxicity in rats. *J Nat Prod*. 2015; 8: 113-122.
3. Mohammed MHH, Hamed ANE, Khalil HE, Kamel MS. Phytochemical and pharmacological studies of *Citharexylum quadrangulare* Jacq leaves. *J Med Plants Res*. 2016; 10: 232-241.
4. Bhutta ZA, Sommerfeld J, Lassi ZS, Salam RA, Das JK. Global burden, distribution, and interventions for infectious diseases of poverty. *Infect Dis Poverty*. 2014; 3: 21.
5. Hotez PJ, Alvarado M, Basáñez MG, Bolliger I, Bourne R, Boussinesq M, et al. The Global Burden of Disease Study 2010: interpretation and implications for the neglected tropical diseases. *PLoS Negl Trop Dis*. 2014; 8: e2865.
6. Dash S, Das C, Sahoo DC, Sahoo AC. Phytochemical composition, anti-inflammatory and analgesic activities of *Tecoma stans* Linn. (Bignoniaceae). *Nat Pharm Technol*. 2011; 1: 5-8.
7. Abdel-Wahab NM, Hamed ANE, Khalil HE, Kamel MS. Pharmacotherapeutic evaluation of *Parmentiera cereifera* Seem. (Family Bignoniaceae) cultivated in Egypt on albino rats. *Eur J Med Plants*. 2015; 8: 29-38.
8. Fischer E, Theisen I, Lohmann LG, Kadereit JW. Bignoniaceae. Flowering plants dicotyledons. *The Families and Genera of Vascular Plants*. K. Kubitzki, Springer Berlin Heidelberg. 2004; 7: 9-13.
9. Sprague TA. *Flora of Tropical Africa*. 1905; 4: 516.
10. Abdel-Hameed UK. Morphological phylogenetics of Bignoniaceae Juss. Beni-Suef Univ J Basic Appl Sci. 2014; 3: 172-177.
11. Lacroix D, Prado S, Deville A, Krief S, Dumontet V, Kasenene J, et al. Hydroperoxy-cycloartane triterpenoids from the leaves of *Markhamia lutea* a plant ingested by wild Chimpanzees. *Phytochemistry*. 2009; 70: 1239-1245.
12. Kernan MR, Amarquaye A, Chen JL, Chan J, Sesin DF, Parkinson N, et al. Antiviral phenyl propanoid glycosides from the medicinal plant *Markhamia lutea*. *J Nat Prod*. 1998; 61: 564-570.
13. Tantangmo F, Lenta BN, BoyomFF, Ngouela S, Kaiser M, Tsamo E, et al. Antiprotozoal activities of some constituents of *Markhamia tomentosa* (Bignoniaceae). *Ann Trop Med Parasitol*. 2010; 104: 391-398.
14. Harborne JB. *Phytochemical Methods a Guide to Modern Techniques of Plant Analysis*. 3rd edn. UK: Chapman & Hall. 1998.
15. Trease GE, Evans WC. *Pharmacognosy*. 14th edn. London: WB Sanders Company. 1985.
16. Makler MT, Hinrichs DJ. Measurement of the lactate-dehydrogenase activity of *Plasmodium falciparum* as an assessment of parasitemia. *Am J Trop Med Hyg*. 1993; 48: 205-210.
17. Mustafa J, Khan SI, Ma GY, Walker LA, Khan IA. Synthesis and anticancer activities of fatty acid analogs of podophyllotoxin. *Lipids*. 2004; 39: 167-172.
18. Ma GY, Khan SI, Jacob MR, Tekwani BL, Li ZQ, Pasco DS, et al. Antimicrobial and antileishmanial activities of hypocrellins A and B Antimicrob Agents Chemother. 2004; 48: 4450-4452.
19. Fournet A, Muñoz V. Natural products as trypanocidal, antileishmanial and antimalarial drugs. *Curr Top Med Chem*. 2002; 2: 1215-1237.
20. Pan L, Lezama-Davila CM, Isaac-Marquez AP, Calomeni EP, Fuchs JR, Satoskar AR, et al. Sterols with antileishmanial activity isolated from the roots of *Pentalinon andrieuxii*. *Phytochemistry*. 2012; 82: 128-135.
21. Ali S, El-Ahmady S, Ayoub N, Singab AN. Phytochemicals of Markhamia Species (Bignoniaceae) and Their Therapeutic Value: A Review. *Eur J Med Plants*. 2015; 6: 124-142.
22. Mazoir N, Benharref A, Bailén M, Reina M, González-Coloma A, Martínez-Díaz RA. Antileishmanial and antitrypanosomal activity of triterpene derivatives from latex of two Euphorbia species. *Z Naturforsch C*. 2011; 66: 360-366.
23. Hoet S, Pieters L, Muccioli GG, Habib-Jiwan J, Opperdoes FR, Quetin-Leclercq J. Antitrypanosomal activity of triterpenoids and sterols from the leaves of *Strychnos spinosa* and related compounds. *J Nat Prod*. 2007; 70: 1360-1363.
24. Desjeux P. Leishmaniasis: current situation and new perspectives. *Comp Immuno I Microbiol Infect Dis*. 2004; 27: 305-318.
25. Newman DJ, Cragg GM. Natural products as sources of new drugs over the last 25 years. *J Nat Prod*. 2007; 70: 461-477.