

# Arrhythmogenic Right Ventricular Cardiomyopathy: From Diagnosis to Treatment

Gregory Dendramis<sup>1\*</sup>, M.D.

<sup>1</sup>Cardiovascular Division, University Hospital "Paolo Giaccone", Palermo, Italy

## Article Information

Received date: Sep 19, 2016

Accepted date: Sep 20, 2016

Published date: Sep 21, 2016

## \*Corresponding author

Gregory Dendramis, C.da S.Caterina, snc, CAP 90037, Piana degli Albanesi, Palermo, Italy, Email: gregorydendramis@libero.it

**Distributed under** Creative Commons CC-BY 4.0

**Keywords** Arrhythmogenic right ventricular cardiomyopathy; Diagnosis; Prognosis; Treatment

## Introduction

Arrhythmogenic Right Ventricular Cardiomyopathy/dysplasia (ARVC) is a genetic form of cardiomyopathy affecting primarily the Right Ventricle (RV) but also may involve the Left Ventricle (LV) and may culminate in life-threatening ventricular arrhythmias, Sudden Cardiac Death (SCD) and/or heart failure. In most cases it is transmitted with an autosomal dominant pattern of inheritance, with incomplete penetrance and variable expression, but there are also some rare autosomal recessive forms as Naxos disease and Carvajal syndrome.

The term "cardiomyopathy" should be preferred to "dysplasia", as already suggested by WHO/International Society and Federation of Cardiology in 1996 and Maron, about classification of cardiomyopathy in 2006 [1,2].

The estimated prevalence of ARVC in the general population is approximately 1:5000, affecting men more frequently than women with a ratio of 3:1 [3,4] and accounts for 11%-22% of cases of SCD in young athletes, resulting in approximately 22% of cases in athletes in northern Italy and about 17% of SCD in young people in the United States [5-7].

The pathologic hallmark of disease is myocardial atrophy (myocyte loss), fibrofatty replacement, fibrosis and ultimate thinning of the wall with chamber dilation and aneurysm formation [8]. The genetics of ARVC support the hypothesis that it may be caused by desmosomal dysfunction (mutation of plakophilin-2, desmoglein-2, desmocollin-2, desmoplakin genes) that lead to impaired mechanical and electrical coupling between individual cells, leading to myocyte uncoupling, especially under conditions that increase myocardial strain, for example during physical effort [9]. These changes may produce electrical instability precipitating ventricular tachycardia and SCD.

Currently only approximately 50% of probands are found to have a pathogenic mutation, possibly due to the high incidence of rare mutations in many patients and actually the most important utility of genetic test are in the screening of asymptomatic family members of probands who have a pathologic genetic abnormality [9].


Diagnosis of ARVC relies on a scoring system, formulated in 2010 by the revisited Task Force, with major and minor criteria based on the demonstration of a combination of defects in RV morphology and function, characteristic depolarization/repolarization electrocardiogram abnormalities (negative T waves and/or "epsilon" wave in right precordial leads), characteristic tissue pathology, typical arrhythmias, family history, and the results of genetic.

Patients are diagnosed as having definite ARVC if they have "4 points" with major criteria equal to 2 points and minor criteria equal to 1 point. Patients whose score totals to "3 points" can be classified as having probable ARVC [9].

Assessment of ventricular structure and function is critical for the diagnosis and prognosis of ARVC. Transthoracic echocardiography provides a noninvasive method of RV imaging, suggestive findings of ARVC include global or segmental wall motion abnormalities (akinesis/dyskinesis) with cavity dilation, hypertrophic RV trabeculation, and systolic dysfunction [9], (Figure), (Table).

To date Cardiac magnetic resonance (MRI) is considered the best imaging modality in evaluating the RV in ARVC and provides tissue characterization and identification of intramyocardial fat and fibrosis (Delayed Enhancement, DE-MRI) in addition to assessment of ventricular structure and function [10,11]. Endomyocardial biopsy has low diagnostic sensitivity for several reasons as the patchy distribution of the disease and the high rate of sampling errors [9].

Symptoms usually appear between the ages of 30-50 and the most common clinical presentation consists of palpitations, syncope and ventricular tachycardia of left bundle branch morphology, especially in young patients [12]. In cases of older patients in whom the disease has been described, the clinical presentation is mainly represented by signs and symptoms of right or biventricular heart failure [13].



**Figure 1:** ARVD: ventriculography (panel A) and cardiac magnetic resonance with areas of dyskinesia of the RV free wall (panel B) and parietal signal alterations (panel C).

**Table 1:** 2010 Task Force Criteria for the Diagnosis of Arrhythmogenic Right Ventricular Cardiomyopathy.

<b>I. Global and/or regional dysfunction and structural alterations</b>	
Major	<ul style="list-style-type: none"> <li>- By 2-dimensional echocardiogram: regional RV akinesia, dyskinesia, or aneurysm and 1 of the following (end diastole):                PLAX RVOT <math>\geq 32</math>mm (corrected for body size [PLAX/BSA] <math>\geq 19</math>mm/m<sup>2</sup>),                PSAX RVOT <math>\geq 36</math>mm (corrected for body size [PSAX/BSA] <math>\geq 21</math>mm/m<sup>2</sup>), Fractional area change <math>\leq 33\%</math>.</li> <li>- By MRI: regional RV akinesia or dyskinesia or dyssynchronous RV contraction and 1 of the following:                Ratio of RV end-diastolic volume to BSA <math>\geq 110</math>mL/m<sup>2</sup> (male) or <math>\geq 100</math>mL/m<sup>2</sup> (female),                RV ejection fraction <math>\leq 40\%</math>.</li> </ul>
Minor	<ul style="list-style-type: none"> <li>- By RV angiography: regional RV akinesia, dyskinesia, or aneurysm.</li> <li>- By 2-dimensional echocardiogram: regional RV akinesia or dyskinesia and 1 of the following (end diastole):                PLAX RVOT <math>\geq 29</math> to <math>&lt; 32</math>mm (corrected for body size [PLAX/BSA] <math>\geq 16</math> to <math>&lt; 19</math> mm/m<sup>2</sup>),                PSAX RVOT <math>\geq 32</math> to <math>&lt; 36</math>mm (corrected for body size [PSAX/BSA] <math>\geq 18</math> to <math>&lt; 21</math> mm/m<sup>2</sup>),                Fractional area change <math>&gt; 33\%</math> to <math>\leq 40\%</math>.</li> <li>- By MRI: regional RV akinesia or dyskinesia or dyssynchronous RV contraction and 1 of the following:                Ratio of RV end-diastolic volume to BSA <math>\geq 100</math> to <math>&lt; 110</math> mL/m<sup>2</sup> (male) or <math>\geq 90</math> to <math>&lt; 100</math> mL/m<sup>2</sup> (female),                RV ejection fraction <math>&gt; 40\%</math> to <math>\leq 45\%</math>.</li> </ul>
<b>II. Tissue characterisation of the wall</b>	
Major	- Residual myocytes $< 60\%$ by morphometric analysis (or $< 50\%$ if estimated), with fibrous replacement of the RV free wall myocardium in $\geq 1$ sample, with or without fatty replacement of tissue on endomyocardial biopsy.
Minor	- Residual myocytes 60% to 75% by morphometric analysis (or 50% to 65% if estimated), with fibrous replacement of the RV free wall myocardium in $\geq 1$ sample, with or without fatty replacement of tissue on endomyocardial biopsy.
<b>III. Repolarisation abnormalities</b>	
Major	- Inverted T waves in right precordial leads (V1, V2, and V3) or beyond in individuals $> 14$ years of age (in the absence of complete RBBB QRS $\geq 120$ ms).
Minor	- Inverted T waves in leads V1 and V2 in individuals $> 14$ years of age (in the absence of complete RBBB) or in V4, V5, or V6/Inverted T waves in leads V1, V2, V3, and V4 in individuals $> 14$ years of age in the presence of complete RBBB.
<b>IV. Depolarisation/conduction abnormalities</b>	
Major	- Epsilon wave (reproducible low-amplitude signals between end of QRS complex to onset of the T wave) in the right precordial leads (V1 to V3).
Minor	- Late potentials by SAECG in $\geq 1$ of 3 parameters in the absence of a QRS duration of $\geq 110$ ms on the standard ECG: filtered QRS duration (fQRS) $\geq 114$ ms; duration of terminal QRS $< 40\mu$ V (low-amplitude signal duration) $\geq 38$ ms; root-mean-square voltage of terminal 40 ms $\leq 20\mu$ V Terminal activation duration of QRS $\geq 55$ ms measured from the nadir of the S wave to the end of the QRS, including R', in V1, V2, or V3, in the absence of complete RBBB.
<b>V. Arrhythmias</b>	
Major	- Nonsustained or sustained ventricular tachycardia of left bundle-branch morphology with superior axis (negative or indeterminate QRS in leads II, III, and aVF and positive in lead aVL).
Minor	- Nonsustained or sustained ventricular tachycardia of RVOT configuration, LBBB morphology with inferior axis (positive QRS in leads II, III, and aVF and negative in lead aVL) or of unknown axis $> 500$ ventricular extrasystoles per 24h (Holter).
<b>VI. Family history</b>	
Major	<ul style="list-style-type: none"> <li>- ARVC confirmed in a first-degree relative who meets current Task Force criteria.</li> <li>- ARVC confirmed pathologically at autopsy or surgery in a first-degree relative.</li> </ul>
Minor	<ul style="list-style-type: none"> <li>- Identification of a pathogenic mutation categorized as associated or probably associated with ARVC in the patient under evaluation.</li> <li>- History of ARVC in a first-degree relative in whom it is not possible or practical to determine whether the family member meets current Task Force criteria.</li> <li>- Premature sudden death (<math>&lt; 35</math> years of age) due to suspected ARVC in a first-degree relative.</li> <li>- ARVC confirmed pathologically or by current Task Force criteria in second-degree relative.</li> </ul>

ARVC, Arrhythmogenic Right Ventricular Cardiomyopathy/dysplasia; aVF, augmented Voltage unipolar left Foot lead; aVL, augmented Voltage unipolar Left arm lead; BSA, Body Surface Area; ECG, Electro Cardio Gram; LBBB, Left Bundle Branch Block; MRI, Magnetic Resonance Imaging; PLAX, Parasternal Long-Axis view; PSAX, Parasternal Short-Axis view; RBBB, Right Bundle Branch Block; RV, Right Ventricular; RVOT, Right Ventricular Outflow Tract; SAECG, Signal-Averaged ECG.

The main goal of therapy is to prevent life-threatening events. To achieve this it is necessary to identify high-risk patients for malignant arrhythmias and SCD that should be considered for ICD implantation in primary prevention. Consequently patients with unexplained syncope, non-sustained VT on noninvasive monitoring, familial history of sudden death, extensive disease including those with LV involvement and good functional status are potential candidates for ICD implantation even in the absence of ventricular arrhythmias. ICD implantation in secondary prevention is obviously mandatory [10,14]. To date VT ablation has an ancillary role and may be helpful in reducing symptoms and ICD firing but not able to prevent SCD [14,15].

Antiarrhythmic drugs, primarily beta blockers, sotalol, and amiodarone have been used for symptomatic control in patients who are not candidates for ICD or as an adjunct therapy to reduce frequent ICD firing due to recurrent VT, but the evidence available has been derived from observational studies [15,20]. Finally, but not least, an important recommendation for patients with ARVC is exercise restriction because an intense physical activity appears to increase the rate of progression of the disease [21,22].

## References

- Richardson P, McKenna W, Bristow M, Maisch B, Mautner B, O'Connell J, et al. Report of the 1995 World Health Organization/International Society and Federation of Cardiology Task Force on the Definition and Classification of cardiomyopathies. *Circulation*. 1996; 93: 841-842.
- Maron BJ, Towbin JA, Thiene G, Antzelevitch C, Corrado D, Arnett D, et al. Contemporary definitions and classification of the cardiomyopathies: an American Heart Association Scientific Statement from the Council on Clinical Cardiology, Heart Failure and Transplantation Committee; Quality of Care and Outcomes Research and Functional Genomics and Translational Biology Interdisciplinary Working Groups; and Council on Epidemiology and Prevention. *Circulation*. 2006; 113: 1807-1816.
- Corrado D, Thiene G. Arrhythmogenic right ventricular cardiomyopathy/dysplasia: clinical impact of molecular genetic studies. *Circulation*. 2006; 113: 1634-1637.
- Gemayel C, Pelliccia A, Thompson PD. Arrhythmogenic right ventricular cardiomyopathy. *J Am Coll Cardiol*. 2001; 38: 1773-1781.
- Dalal D, Nasir K, Bomma C, Prakasa K, Tandri H, Piccini J, et al. Arrhythmogenic right ventricular dysplasia: a United States experience. *Circulation*. 2005; 112: 3823-3832.
- Corrado D, Fontaine G, Marcus FI, McKenna WJ, Nava A, Thiene G, et al. Arrhythmogenic right ventricular dysplasia/cardiomyopathy: need for an international registry. Study Group on Arrhythmogenic Right Ventricular Dysplasia/Cardiomyopathy of the Working Groups on Myocardial and Pericardial Disease and Arrhythmias of the European Society of Cardiology and of the Scientific Council on Cardiomyopathies of the World Heart Federation. *Circulation*. 2000; 101: E101-106.
- Corrado D, Basso C, Schiavon M, Thiene G. Screening for hypertrophic cardiomyopathy in young athletes. *N Engl J Med*. 1998; 339: 364-369.
- Thiene G, Basso C, Danieli G, Rampazzo A, Corrado D, Nava A. Arrhythmogenic right ventricular cardiomyopathy a still underrecognized clinic entity. *Trends Cardiovasc Med*. 1997; 7: 84-90.
- Calkins H. Arrhythmogenic right ventricular dysplasia. *Curr Probl Cardiol*. 2013; 38: 207-208.
- Jain A, Tandri H, Calkins H, Bluemke DA. Role of cardiovascular magnetic resonance imaging in arrhythmogenic right ventricular dysplasia. *J Cardiovasc Magn Reson*. 2008; 20: 10-32.
- Tandri H, Saranathan M, Rodriguez ER, Martinez C, Bomma C, Nasir K, et al. Noninvasive detection of myocardial fibrosis in arrhythmogenic right ventricular cardiomyopathy using delayed-enhancement magnetic resonance imaging. *J Am Coll Cardiol*. 2005; 45: 98-103.
- Romero J, Mejia-Lopez E, Manrique C, Lucariello R. Arrhythmogenic Right Ventricular Cardiomyopathy (ARVC/D): A Systematic Literature Review. *Clinical Medicine Insights: Cardiology*. 2013; 7: 97-114.
- More D, O'Brien K, Shaw J. Arrhythmogenic right ventricular dysplasia in the elderly. *Pacing Clin Electrophysiol*. 2002; 25: 1266-1269.
- Epstein AE, DiMarco JP, Ellenbogen KA, Estes NA 3<sup>rd</sup>, Freedman RA, Gettes LS, et al. ACC/AHA/HRS 2008 Guidelines for Device-Based Therapy of Cardiac Rhythm Abnormalities: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the ACC/AHA/NASPE 2002 Guideline Update for Implantation of Cardiac Pacemakers and Antiarrhythmia Devices): developed in collaboration with the American Association for Thoracic Surgery and Society of Thoracic Surgeons. *Circulation*. 2008; 117: e350-408.
- Zipes DP, Camm AJ, Borggrefe M, Buxton AE, Chaitman B, Fromer M, et al. ACC/AHA/ESC 2006 guidelines for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: a report of the American College of Cardiology/American Heart Association Task Force and the European Society of Cardiology Committee for Practice Guidelines (Writing Committee to Develop Guidelines for Management of Patients With Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death). *J Am Coll Cardiol*. 2006; 48: e247-346.
- Corrado D, Calkins H, Link MS, Leoni L, Favale S, Bevilacqua M, et al. Prophylactic implantable defibrillator in patients with arrhythmogenic right ventricular cardiomyopathy/dysplasia and no prior ventricular fibrillation or sustained ventricular tachycardia. *Circulation*. 2010; 122: 1144-1152.
- Hodgkinson KA, Parfrey PS, Bassett AS, Kupprion C, Drenkhahn J, Norman MW, et al. The impact of implantable cardioverter-defibrillator therapy on survival in autosomal-dominant arrhythmogenic right ventricular cardiomyopathy (ARVD5). *J Am Coll Cardiol*. 2005; 45: 400-408.
- Zwanziger J, Hall WJ, Dick AW, Zhao H, Mushlin AL, Hahn RM, et al. The cost effectiveness of implantable cardioverter-defibrillators: results from the Multicenter Automatic Defibrillator Implantation Trial (MADIT)-II. *J Am Coll Cardiol*. 2006; 47: 2310-2318.
- Corrado D, Leoni L, Link MS, Della Bella P, Gaita F, Curnis A, et al. Implantable cardioverter-defibrillator therapy for prevention of sudden death in patients with arrhythmogenic right ventricular cardiomyopathy/dysplasia. *Circulation*. 2003; 108: 3084-3091.
- Steinberg JS, Martins J, Sadanandan S, Goldner B, Menchavez E, Domanski M, et al. Antiarrhythmic drug use in the implantable defibrillator arm of the Antiarrhythmics Versus Implantable Defibrillators (AVID) Study. *Am Heart J*. 2001; 142: 520-529.
- Kirchhof P, Fabritz L, Zweiner M, Witt H, Schafers M, Zellerhoff S, et al. Age and training dependent development of arrhythmogenic right ventricular cardiomyopathy in heterozygous plakoglobin deficient mice. *Circulation*. 2006; 114: 1799-1806.
- Dendramis G, Orlando D, Paleologo C, Ciaramitaro G. A case of arrhythmogenic right ventricular cardiomyopathy without ventricular arrhythmias, *Journal of Indian College of Cardiology*. 2014; 4: 124-127.